

X-ray bursts from 1RXS J170930.2-263927 = XTE J1709-267

M. Cocchi¹, A. Bazzano, L. Natalucci and P. Ubertini

Istituto di Astrofisica Spaziale (*IAS/CNR*), via Fosso del Cavaliere, 00133 Roma, Italy

J. Heise, J.M. Muller², M.J.S. Smith² and J.J.M. in 't Zand

Space Research Organization Netherlands (*SRON*), Sorbonnelaan 2, 3584 TA Utrecht, The Netherlands

ABSTRACT

We report evidence of type-I X-ray bursting activity from the transient *1RXS J170930.2-263927* when it was in outburst early 1997. This identifies the source as a probable low-mass X-ray binary containing a neutron star. The error boxes of the detected bursts and of the persistent emission, as obtained with the *Wide Field Cameras* on board the *BeppoSAX* satellite, rule out an association with the proposed radio counterpart (Hjellming & Rupen 1997).

Subject headings: binaries: close, individual (*1RXS J170930.2-263927*) — X-rays: bursts

1. Introduction

1RXS J170930.2-263927 was first observed in hard X-rays with the *Proportional Counter Array (PCA)* on board the *Rossi X-ray Timing Explorer (RXTE)* (Marshall et al. 1997). The source, reported as a new *XTE* transient source (*XTE J1709-267*), was detected for about 20 s during two different satellite manoeuvres, on 1997 January 16.016 and 1997 January 18.147 respectively. The uncertainty of the source position was estimated to be 6', suggesting a possible association of *XTE J1709-267* with the soft X-ray source *1RXS J170930.2-263927* included in the *ROSAT All-Sky Survey* catalogue (Voges et al. 1996), 5' from the *PCA* centroid position (see Figure 3).

A possible radio counterpart of the source was found in the 1.42 and 4.9 GHz bands (Hjellming & Rupen 1997). The radio position, given with better than 1'' uncertainty, is 2'8 from the *PCA* centroid position and 6' from the *ROSAT* position. So the hard X-ray source could only be associated either with *1RXS J170930.2-263927* or with the proposed radio counterpart, but not

¹e-mail address: wood@ias.rm.cnr.it

²also *BeppoSAX* Science Data Centre, Nuova Telespazio, via Corcolle 19, 00131 Roma, Italy

with both.

The X-ray characteristics of the *RXTE* source are not known in detail. The spectrum obtained by the *PCA* could be fitted with a power law function³ with a 3.06 ± 0.06 photon index or by a thermal bremsstrahlung model with $kT = 4.7 \pm 0.2$ keV. In both cases the column density was found in excess of $2 \times 10^{22} \text{ cm}^{-2}$, thus indicating the source to be close to the Galactic Center. During the *PCA* observations the source had an intensity of 140 mCrab in the 2-10 keV band, while the 1-2.4 keV flux of *1RXS J170930.2-263927* at its detection time (spring 1996) was only 13 mCrab.

After its detection in hard X-rays, *1RXS J170930.2-263927* was monitored by the *All Sky Monitor (ASM)* on board *RXTE* in the 2-10 keV range. The source was still bright (140-200 mCrab) up to February 20, then declined to 80-120 mCrab (March 12-18) and to 25-65 mCrab since March 19. The source was no longer in outburst since April 7, its daily averaged flux being below 13 mCrab. The *RXTE-ASM* lightcurve of the outburst is displayed in Figure 1. The lightcurve does not show uncommon features when compared with the average lightcurves of the X-ray novae. Following the morphological classification proposed by Chen, Shrader, & Livio 1997, we can associate the source with the fast-rise exponential-decay class, even if the decay looks almost linear and perhaps dropping faster about 50 days after the maximum. The *ASM* peak luminosity of the outburst is 0.21 Crab, a rather low but not uncommon value among the X-ray novae, where selection effects in the studied samples are not negligible. We can derive that at the outburst epoch the source intensity increased by more than an order of magnitude.

In the next Section we summarize the *BeppoSAX-WFC* telescope characteristics and report on the observations of *1RXS J170930.2-263927*. In Section 3 the performed spectral and time analysis are presented, while in Section 4 the implications of our results on the knowledge of the source characteristics are briefly discussed. In particular, the association of *1RXS J170930.2-263927* with the class of transient neutron-star low-mass X-ray binaries is proposed.

2. Observations

The *Wide Field Cameras (WFC)* on board the *BeppoSAX* satellite consist of two identical coded mask telescopes (Jager et al. 1997). The two cameras point in opposite directions each covering a $40^\circ \times 40^\circ$ field of view. This field of view is the largest ever flown for an X-ray imaging device. With its source location accuracy better than 0'6 (68% confidence level), a time resolution of 0.244 ms, and an energy resolution of 18% at 6 keV, the *WFCs* are very effective in studying hard X-ray (2-28 keV) transient phenomena. The imaging capability, combined with the good instrument sensitivity (a few mCrab in 10^4 s), allows an accurate monitoring of complex sky regions, like the Galactic Bulge.

The data of the two cameras are systematically searched for bursts and flares by analyzing the time

³ i.e. $n(E) \propto E^{-\Gamma}$, where $n(E)$ is the number of photons with energy E and Γ is the photon index.

profiles of the detectors in the 2-11 keV energy range with a 1 s time resolution. Reconstructed sky images are generated for any statistically meaningful enhancement, to identify possible bursters. The accuracy of the reconstructed position, which of course depends on the burst intensity, is typically better than 5'. This analysis procedure demonstrated its effectiveness throughout the Galactic Bulge *WFC* monitoring (Cocchi et al. 1997a), leading to the identification of ~ 530 X-ray bursts, 156 of which from the *Bursting Pulsar GRO J1744-28*, in a total of about 1100 ks observing time (the exposure time values are corrected for all dead times due to Earth occultation and South Atlantic Anomaly). A total of 12 new X-ray bursting sources were found.

1RXS J170930.2-263927 is in the the field of view whenever the *WFCs* point at the Galactic Centre region, being 9.4 away from the Sgr A position. The Galactic Bulge field was observed for a total of about 1100 ks, the monitoring being divided in 4 observation campaigns during 1996 through 1998. Unfortunately, *1RXS J170930.2-263927* was out of visibility at the epoch of the outburst onset (January '97), so only data from the declining phase and quiescence are available.

During a 102.6 ks observation started on 1997 March 18.04 a burst was detected at a position consistent with that of *1RXS J170930.2-263927* at March 18.21058 *UT* (Cocchi et al. 1997b). The event occurred during the egress from the Earth's occultation. Two more bursts were detected from the same position in a different observation shift that started on April 13.16 and lasting for 97.8 ks. The time profiles of the three bursts are plotted in Figure 2. The displayed profiles of the bursts are detector profiles constructed by integrating the photons in the source-illuminated part of the detector. This improves the signal to noise ratio of the event. The background is the sum of the diffuse X-ray background, the particles background and the contamination of other sources in the field of view. Source contamination is the dominating background component for crowded fields like the Galactic Bulge. Nevertheless, the probability of source confusion during a short time scale event like an X-ray burst is negligible. Details on the three observed bursts are summarized in Table 1.

Burst searches from *1RXS J170930.2-263927* were performed on all the data available from the '96-'98 *BeppoSAX-WFC* Galactic Bulge monitoring campaigns but no other events were detected.

3. Data Analysis

The steady emission of *1RXS J170930.2-263927* during its declining phase in March '97 has been investigated. We analyzed data from two observations which started on March 13.52 and March 18.04 (U.T.) for a total of 170.6 ks exposure time.

The intensity of the source was found to be fairly constant throughout the two observations, the average flux being about 28 mCrab. The data was split in five data subsets in order to investigate possible spectral variability on 1 day timescale. The fitted parameters over all the subsets are consistent with a constant spectrum, thus excluding spectral variability during the involved epochs. Two spectral models were applied to the average spectrum over all the five subsets: an absorbed power law function and a thermal bremsstrahlung spectrum. The calculated parameters,

which are consistent with the ones reported by *RXTE-PCA*, are shown in Table 2.

An absorbed blackbody model was adopted to fit the spectra of the three bursts. To better constrain the fit, the N_{H} parameter was kept fixed ($N_{\text{H}} = 2 \times 10^{22} \text{ cm}^{-2}$) according to what was obtained from the spectral analysis of the steady emission by using *WFC* and *RXTE-PCA* data. Moreover, two time-resolved spectra were obtained for the 2nd burst, i.e. a "peak spectrum" (data from the first 6 s of the burst) and a "tail spectrum" (the next 9 s of burst data). A summary of the spectral parameters of the three bursts is given in Table 1.

Blackbody spectra allow to determine the relationship between the average radius of the emitting sphere R_{km} (in units of km) and the source distance $d_{10 \text{ kpc}}$ (in units of 10 kpc). In Table 1 the values of the $R_{\text{km}}/d_{10 \text{ kpc}}$ ratios are given, assuming isotropic emission and not correcting for gravitational redshift and conversion to true blackbody temperature from color temperature (see Lewin, van Paradijs, & Taam 1993 for details).

The calculated *WFC* centroid position of the observed hard X-ray source was found to be consistent with that of the soft *ROSAT* source *1RXS J170930.2-263927* (Cocchi et al. 1997a). In Figure 3 a sky map is shown with the error circles of the position we calculated for the March 18 persistent emission and for the three bursts. The *WFC* position is almost coincident with the *ROSAT* one, thus associating the hard X-ray *XTE* source with the soft X-ray *ROSAT* source *1RXS J170930.2-263927*. A possible association of both the soft and the hard X-ray sources with the proposed radio counterpart, which lies more than 5' away, is ruled out.

4. Discussion

Up to date, transient phenomena like hard X-ray bursts with typical durations less than a few minutes are classified in two main types (Hoffman, Marshall, & Lewin 1978).

Type-I bursts originate from thermonuclear flashes onto a neutron star surface (Woosley & Taam 1976, Joss 1978, Taam & Picklum 1979). All the type-I bursting sources are associated with low mass binary systems (LMXBs), following the identification of their optical counterparts or, indirectly, from spectral or stellar population (e.g. globular cluster sources) characteristics (van Paradijs 1995). The observed spectra of the bursts are typically well fitted by thermal blackbody emission models having temperatures $kT \lesssim 3 \text{ keV}$ (Swank et al. 1977). The time profiles of type-I bursts show fast rise times, ranging from less than 1 s up to 10 s, and longer decay times, ranging from seconds to minutes. They strongly depend on photon energy, decays being shorter at higher energies, so that bursts appear to soften during the decay. This softening has been suggested to originate from the cooling of the neutron star photosphere (e.g. Lewin, van Paradijs, & Taam 1995).

Type-II bursts have been observed only in two peculiar cases, namely the *Rapid Burster MXB 1730-335* (Lewin et al. 1976) and the *Bursting Pulsar GRO J1744-28* (Kouveliotou et al. 1996). They show no spectral softening and are frequently repeated in time, intervals ranging from a few seconds to hours. Blackbody fits to the type-II bursts are unacceptably poor. Comptonized

models for *MXB 1730-335* (Stella et al. 1988) and a power law function model for *GRO J1744-28* (Giles et al. 1996) fit the data better. Moreover, the *Bursting Pulsar* spectra of the steady emission and the bursts are very similar. Type-II bursts are thought to originate from accretion instabilities (Lewin et al. 1995).

On the basis of their spectral properties and time profiles, we interpret the three bursts detected from *1RXS J170930.2-263927* as type-I bursts. Even though the statistical quality of the data is limited, the blackbody emission and the measured color temperatures are consistent with type-I bursts. Spectral softening is marginally observed in time resolved spectra of burst 2 and energy resolved time profiles of all the bursts (see Table 1). Average decay times of 8 ± 2 s (2-6 keV) and 4 ± 1 s (6-18 keV) are obtained.

Type-I bursts strongly suggest a neutron star nature for the binary system. This indicates *1RXS J170930.2-263927* as a transient neutron-star LMXB.

In principle, it is possible that *1RXS J170930.2-263927* had bursting behaviour even before the January 1997 outburst, as the bursts occurrence doesn't seem to be related to the persistent flux (bursts 2 and 3 occurred when the steady flux had dropped by a factor $\gtrsim 10$ with respect to the time of burst 1, see Table 1). On the other hand, it is possible that the outburst itself triggered the type-I burst mechanism, due to an increased accretion rate from the companion star and that the physical conditions that activated the bursts persisted even after the source entered a quiescent phase. This scenario seems likely since no bursting activity was detected from *1RXS J170930.2-263927* in the autumn '96 and autumn '97 observation campaigns.

Bursting activity from LMXB transients have already been reported in about 10 cases (e.g. *Rapid Burster*, *Aql X-1*, *Cen X-4*, *0748-673*, *1658-298*, *SAXJ 1808.4-3658*, see Hoffman et al. 1978, Tanaka & Shibazaki 1996, Lewin et al. 1995, and references therein, In 't Zand et al. 1998), thus indicating the sources to be neutron-star binaries. Most ($\simeq 85\%$) of the LMXB harbouring a neutron star are persistent sources, while all the black hole candidates in low-mass systems are transient sources (van Paradijs 1995). Among the LMXB transients a quite large fraction of sources ($\simeq 30\%$, according to Chen et al. 1997, $\simeq 45\%$, according to Tanaka & Shibazaki 1996) are neutron-star systems, the rest being black hole binaries. The population of black hole LMXB could be overestimated, since most of them are suggested as black hole candidates on the basis of their spectral characteristics only. Actually, for only 7 out of about 40 known transient LMXB the available mass functions suggest black hole systems (Chen et al. 1997).

Thus, the detection of type-I X-ray burst in X-ray novae can help to better constrain the populations.

Due to the uncertainty in the distance of the source, interesting parameters such as the luminosity of *1RXS J170930.2-263927* both in outburst and in quiescence, the blackbody emitting region radius during the bursts and the peak luminosities of the bursts cannot be satisfactorily constrained. The interpolated N_{H} value computed at the *ROSAT* given position is $2.3 \times 10^{21} \text{ cm}^{-2}$ (Dickey & Lockman 1990), about an order of magnitude lower than the measured value, thus suggesting intrinsic absorption. This makes it difficult to use the observed N_{H} as a distance

indicator.

A rough upper limit for the source distance can be obtained assuming the observed bursts had luminosities below the Eddington limit ($L_{\text{Edd}} \simeq 2 \times 10^{38} \text{ erg s}^{-1}$). Low blackbody temperatures ($< 2.5 \text{ keV}$) and no evidence of double-peaked profiles (a clue of photospheric radius expansion) support this assumption for type-I bursts (Lewin et al. 1995). Correcting for interstellar absorption, for burst 2 we obtain a bolometric flux of $L_{\text{b}} = 1.6 \times 10^{-8} \text{ erg s}^{-1} \text{ cm}^{-2}$. Assuming $L_{\text{b}} \leq L_{\text{Edd}}$, this leads to a maximum distance of $10 \pm 1 \text{ kpc}$, but a lower distance value is likely since the source is at relatively high (7.9°) galactic latitude. We then derive upper limits of $\sim 8 \times 10^{37} \text{ erg s}^{-1}$ for the outburst peak luminosity and $\sim 1 \times 10^{36} \text{ erg s}^{-1}$ for the persistent emission at the bursts 2,3 epochs. An upper limit of $\sim 10 \text{ km}$ is also obtained for the average radius of the blackbody emitting region during the observed bursts, a value supporting the neutron star identification.

We thank the staff of the *BeppoSAX Science Operation Centre* and *Science Data Centre* for their help in carrying out and processing the *WFC* Galactic Centre observations. The *BeppoSAX* satellite is a joint Italian and Dutch program. M.C., A.B., L.N. and P.U. thank Agenzia Spaziale Nazionale (*ASI*) for grant support.

REFERENCES

- Chen, W., Shrader, C.R., & Livio, M. 1997, ApJ, 491, 312.
- Cocchi, M., et al. 1997a, in *"The Active X-ray Sky: results from BeppoSAX and RossiXTE"*, ed. L. Scarsi, H. Bradt, P. Giommi, & F. Fiore, Nucl. Phys. B Proc. Suppl., in press
- Cocchi, M., et al. 1997b, in *"The Active X-ray Sky: results from BeppoSAX and RossiXTE"*, ed. L. Scarsi, H. Bradt, P. Giommi, & F. Fiore, Nucl. Phys. B Proc. Suppl., in press
- Dickey, J.M., & Lockman, F.J. 1990, A&A Rev., 28, 215.
- Giles, A.B., et al. 1996, ApJ, 469, L25.
- Hjellming, R.M., & Rupen, M.P. 1997, IAUC 6547.
- Hoffman, J.A., Marshall, H.L., & Lewin, W.H.G. 1978, Nature, 271, 630.
- In 't Zand, J.J.M., et al. 1998, A&A, 331, L25.
- Jager, R., et al. 1997, A&A, 125, 557.
- Joss, P.C. 1978, ApJ, 225, L123.
- Kouveliotou, C., et al. 1996, Nature, 379, 799.
- Lewin, W.H.G., van Paradijs, J., & Taam, R.E. 1993, Space Sci. Rev., 62, 223.
- Lewin, W.H.G., et al. 1976, ApJ, 207, L95.
- Lewin, W.H.G., van Paradijs, J., & Taam, R.E. 1995, in *"X-ray Binaries"*, ed. W. Lewin, J. van Paradijs, & E. van den Heuvel, Cambridge University Press, Cambridge, p. 175
- Marshall, F.E., Swank, J.H., Thomas, B., Angelini, L., Valinia, A., & Ebisawa, K. 1997, IAUC 6543.
- Stella, L., Haberl, F., Lewin, W.H.G., Parmar, A.N., van Paradijs, J., & White, N.E. 1988, ApJ, 324, 379.
- Swank, J.H., Becker R.H., Boldt, E.A., Holt, S.S., Pravdo, S.H., & Serlemitsos, P.J. 1977, ApJ, 212, L73.
- Taam, R.E., & Picklum, R.E. 1979, ApJ, 233, 327.
- Tanaka, Y., & Shibazaki, N. 1996, *"X-ray novae"*, Annu.Rev.Astron.Astrophys., 34, 607.
- van Paradijs, J. 1995, in *"X-ray Binaries"*, ed. W. Lewin, J. van Paradijs, & E. van den Heuvel, Cambridge University Press, Cambridge, p. 536

Voges, W., et al. 1996, IAUC 6420.

Woosley, S.E., & Taam, R.E. 1976, *Nature*, 263, 101.

Table 1. Summary of the *XTE J1709-267* bursts characteristics

	burst 1	burst 2	burst 3
Burst time	March 18.21058	April 13.52963	April 14.59554
Burst e -folding time (2-11 keV)	5.2 ± 1.8 s	7.2 ± 2.8 s	10.0 ± 3.5 s
Maximum burst intensity flux (2-11 keV)	0.48 ± 0.13 Crab	0.50 ± 0.12 Crab	0.48 ± 0.13 Crab
Steady intensity flux (2-10 keV)	26 ± 1 mCrab	< 3 mCrab ^a	< 3 mCrab ^a
Burst e -folding time (2-6 keV)	6.6 ± 3.2	7.5 ± 3.5 s	11.9 ± 4.8 s
Burst e -folding time (6-18 keV)	2.6 ± 1.4	5.5 ± 2.8 s	7.1 ± 4.2 s
Blackbody temperature (keV)	2.10 ± 0.16	1.79 ± 0.16	1.73 ± 0.16
Reduced χ^2 ^b	0.93	0.65	0.91
<i>peak</i> blackbody kT (keV)	–	1.83 ± 0.26	–
<i>tail</i> blackbody kT (keV)	–	1.69 ± 0.28	–
$R_{\text{km}}/d_{10 \text{ kpc}}$	7.9 ± 2.3	11.0 ± 3.3	11.6 ± 3.7

^a 3σ upper limit

^b26 d.o.f.

Table 2. Summary of the persistent emission characteristics

Model	Model parameter	N_{H} (10^{22} cm^{-2})	reduced χ^2 ^b
BR ^a :	$kT = 5.77 \pm 0.63 \text{ keV}$	0.148 ± 0.453	0.94
PL ^a :	$\Gamma = 2.67 \pm 0.13$	2.27 ± 0.63	1.02

^aBR = bremsstrahlung, PL = power law

^b144 d.o.f.




Fig. 1.— *RXTE*-*ASM* lightcurve of *1RXS J170930.2-263927*. The markers 1,2,3 indicate the epochs of the observed *WFC* bursts.




Fig. 2.— 2–11 keV time profiles of the observed bursts.




Fig. 3.— A sky map of relevant error regions. Small solid circle: *1RXS J170930.2-263927* error circle (Voges et al. 1996). Large solid circle: *WFC-SAX* steady source error circle (99% confidence). Dotted circles: error circles of the three *WFC*-detected bursts (99% confidence). Large dashed circle: *XTE J1709-267* error circle (only systematic errors, Marshall et al. 1997). Asterisk: position of the suspected radio counterpart (Hjellming & Rupen 1997).